

# Medications to Avoid Before Surgery

The following drugs contain aspirin or other products which can cause increased bleeding during surgery and should not be taken for 10 days prior to surgery. For relief of minor pain, you MAY take Tylenol (acetaminophen).

Advil	Doan's Regular and	Nabumeton
Aggrenox	Extra Strength	Naprosyn
Aleve	Dolobid	Naprosyn Pepto Bismol
Alka Seltzer	Dristan	Tablets and Liquid
Alcohol	Duradyne Tablets	Naproxen
Anacin	Easprin	Norgesic Forte
Anaprox	Ecotrin	Omega-3
Ansaid	Eliquis	Orudis
Arthotec	Empirin	Oruvail
Aspirin	Enbrel	Oxaprozin
APC	Equagesic Tablets	Percodan
Apixaban	Etodolac	Persantine
BC Tablets or Powders	Excedrin	Piroxicam
BC Cold Powder	Feldene	Plavix
Brilinta	Fiorinal	Pradaxa
Brufen	Fish Oil	Quagesic
Bufferin	Flurbiprofen Sodium	Relafen
Cama Arthritis Pain	4 Way Cold Tablets	Robaxisal
Reliever	Goody's Headache	Rufen
Cataflam	Powder or Tablets	Sine Aid
Celebrex	Ibuprofen	Soma Compound
Cephalgesics	Indomethacin	Sulindac
Cilostazol	Indocin	Trandate
Clinoril	Ketoprofen	Trental
Cogesprin	Lodine	Trilisate
Coricidin	Lovenox	Vanquish
Coumadin	Meclomen	Vitamin E
Dabigatran	Medipren	Voltaren
Darvon	Meloxicam	Warfarin
Darvon with Aspirin	Midol 200	Wesprin
Daypro	Midol PMS Caplets	Xarelto
Diclofenac	Mobic	Zavtrin
Diflunisal	Motrin	Zoprin
Disalcid	Napreelan	<b>Herbal Supplements</b>

Before stopping any of these medications, be sure to consult the physician who ordered them. Some, such as Coumadin (Warfarin) and Plavix, are ordered to prevent or treat serious conditions such as deep venous thrombosis, pulmonary embolisms, and other heart problems. This is not an all-inclusive list. If you are unsure if you are taking an aspirin product or an anti-inflammatory, please ask your doctor, nurse or pharmacist.