IBUPROFEN DOSING CHART

WEIGHT	DOSE	INFANT DROPS	CHILDREN'S Liquid	CHILDREN'S CHEWS	JRSTRENGTH TABS	ADULT Regular-strengthtabs
11–16 lbs	50 mg	1.25 mL	2.5 mL	<u> </u>	—	_
17–22 lbs	75 mg	1.875 mL	3.75 mL	_		_
23–33 lbs	100 mg	2.5 mL	5 m L			
34-44 lbs	150 mg	3.75 mL	7.5 mL	3 chews		
45–55 lbs	200 mg	5 m L	10 mL	4 chews	2 tabs	
56–66 lbs	250 mg		12.5 mL	5 chews	2.5 tabs	1 tab
67–77 lbs	300 mg	_	15 mL	6 chews	3 tabs	1–1.5 tabs
78–88 lbs	350 mg	_	17.5 mL	7 chews	3.5 tabs	1.5 tabs
89+ lbs	400 mg	-	20 mL	8 chews	4 tabs	2 tabs

5 SAFETY TIPS FOR IBUPROFEN

- I. Don't give it to a baby under 6 months without a doctor's approval.
- 2. Always use the measuring device that comes with the medicine — not a spoon from the kitchen.
- 3. Don't confuse infant drops with children's liquid. Infant drops are much more concentrated.
- 4. The proper dosage for your child is based on weight, not age. If you don't know how much your child weighs, and he's too young to stand on a scale himself, weigh yourself while holding him, and then weigh yourself alone. Subtract your weight from the combined weight to get his current weight.
- 5. Don't exceed 4 doses in a 24 hour period.

ACTIVE INGREDIENTS

- Infant drops: 50 mg /1.25 mL) per dropper (ie. 75mg/1.875 mL syringe)
- Note: The drops come with either a dropper or a syringe, depending on the brand.
- Children's liquid: 100 mg per teaspoon (5 mL) (shake well before using)
- Children's chewables: 50 mg each
- Junior-strength chewables or caplets: 100 mg each
- Adult regular-strength tablets: 200 mg each