Gastroenterology Adam Niemann, MD



1118 Hampshire St, Second Floor Quincy, IL 62301

DURING CAPSULE ENDOSCOPY - PATIENT INSTRUCTIONS

You have just swallowed a capsule endoscope. This sheet contains information about what to expect over the next 8 hours. Please call our office if you have severe or persistent abdominal or chest pain, fever, difficulty swallowing, or if you have any questions. Our phone number is (217) 222-6550, option #1, extension 6407.

TIME OF CAPSULE INGESTION =

- 1. You may drink clear liquids (water, apple juice) 2 hours after swallowing the capsule. You may also take your medications 4 hours after swallowing capsule.
- 2. Do not exercise. Avoid heavy lifting. You may walk, sit and lie down. You can drive a car. You may return to work if your work allows avoiding unsuitable environments and/or physical movements.
- 3. Avoid going near MRI machines and radio transmitters. You may use a computer, radio, stereo, or cell phone.
- 4. Do not stand directly next to another patient undergoing capsule endoscopy.
- 5. Try not to touch the recorder or the sensor array. Do not remove the sensor array.
- 6. Avoid getting the data recorder or sensor array wet.
- 7. Observe the LED light on the data recorder at least every 15 minutes. If the light stops blinking, document the time and call our office.

Return to the office at ______ to have the equipment removed.