

COLONOSCOPY INSTRUCTIONS SHEET/GOLYTELY – TWO DAY

To schedule your colonoscopy or if you have any questions about your procedure, call Dr. Niemann's office at (217) 222-6550, option #1, extension 3636.

Your appointment for your colonoscopy is _____ and arrival time is _____ at the Quincy Medical Group Surgery Center at 3301 Broadway, Second Floor, Quincy, IL 62301.

TIME: Quincy Medical Group Clinical Team will call you with the arrival time one day prior to your procedure. If you need to cancel the day of the procedure, call the Quincy Medical Group Surgery Center at (217) 277-4090. **DO NOT CALL THIS NUMBER TO SCHEDULE THE APPOINTMENT.**

You will need to pick up 2 gallons of GoLYTELY Prep solution. A prescription has been sent to your pharmacy.

1. **DAY ONE:** Be on a full-liquid diet all day long _____. Full liquids allowed are: milk, milk products including non-dairy creamers, coffee (black), tea, water, clear fruit juices (grape, apple, cranberry), bouillon, jello, soda, popsicles, or Gatorade. **NO RED/BLUE COLORING.** No solid food. No food with seeds or corn 3 days prior to the procedure.
2. Start GoLYTELY Prep at 4 p.m. _____. Drink one an 8-ounce glass every 15-20 minutes until finished. You may place the solution on ice – but no other additives.
3. **DAY TWO:** Be on a clear-liquid diet all day _____. Clear liquids are anything you can pour into a glass and see through. Liquids allowed are: coffee (black), tea, water, clear fruit juices (grape, apple, cranberry), bouillon, jello, soda, popsicles and Gatorade. **NO RED/BLUE COLORING.** No solid food, milk or milk products including non-dairy creamers.
4. Start GoLYTELY Prep at 4 p.m. _____. Drink one 8-ounce glass every 15-20 minutes until finished. You may place the solution on ice – but not other additives.

WHAT TO EXPECT: After 3 hours of drinking prep, stools should be watery and you should be able to see through toilet water. If not, call Physician Exchange at (217) 222-2088.

5. You may continue with clear liquids until 4 hours prior to the procedure.
6. Since you will have a mild sedative the day of the procedure, you must be accompanied by someone who can drive you home after your procedure.
7. You must **DISCONTINUE** the following medications 7 days prior to your procedure: aspirin, Motrin, ibuprofen, Nuprin, Advil, Aleve, any type of arthritis medication, nonsteroidal anti-inflammatory drugs (NSAIDS), iron supplements, vitamins that contain iron, Aggrenox, any blood-thinning medications such as Plavix or Ticlid, and all herbal supplements.
8. Discontinue Coumadin 3 days prior.
9. Please be sure to take any heart or blood pressure medications with sips of water the morning of procedure.
10. You may take Tylenol products or Celebrex.

IF YOU ARE DIABETIC: The day prior to the procedure you may take your oral medication. If on insulin, take only ½ dose of insulin. Do not take oral diabetic medications on the day of the procedure. Only take ½ dose of insulin the morning of the procedure. If taking Byetta, Januvia or Victoza, hold day before procedure and morning of procedure.