

TYLENOL™ DOSING CHART

WEIGHT	DOSE	OLD INFANT DROPS	NEW INFANT LIQUID	CHILDREN'S LIQUID	CHILDREN'S CHEWS/MELTAWAYS	JR.-STRENGTH CHEWS/MELTAWAYS	ADULT REG.-STRENGTH TABS	ADULT EXTRA-STRENGTH TABS
6–11 lbs	40 mg	0.4 mL in dropper	1.25 mL in syringe	—	—	—	—	—
12–17 lbs	80 mg	0.8 mL in dropper	2.5 mL in syringe	2.5 mL in cup	—	—	—	—
18–23 lbs	120 mg	1.2 mL in dropper	3.75 mL in syringe	3.75 mL in cup	1.5 tab	—	—	—
24–35 lbs	160 mg	1.6 mL in dropper	5 mL in syringe	1 tsp (5 mL in cup)	2 tabs	—	—	—
36–47 lbs	240 mg	—	—	7.5 mL in cup	3 tabs	1.5 tabs	—	—
48–59 lbs	320 mg	—	—	10 mL in cup	4 tabs	2 tabs	1 tab (325 mg)	—
72–95 lbs	480 mg	—	—	15 mL in cup	6 tabs	3 tabs	1–1.5 tabs	—
96+ lbs	640 mg	—	—	20 mL in cup	8 tabs	4 tabs	2 tabs	1 tab (500 mg)

FIVE SAFETY TIPS FOR ACETAMINOPHEN

1. Don't give to a baby under 3 months old without a doctor's approval.
2. Don't confuse infant drops with the new infant liquid. Infant drops are three times more concentrated. Learn more at www.babycenter.com/acetaminophen.
3. Always use the measuring device that comes with the medicine.
4. The proper dosage is based on weight, not age. To determine the weight of a very young child, weigh yourself and then weigh yourself while holding your child. Then subtract your weight from the combined weight.
5. Don't exceed five doses in 24 hours.

ACTIVE INGREDIENTS

OLD INFANT DROPS: 80 mg/0.8 mL per dropper (shake well before using)

NEW INFANT LIQUID: 160 mg/5 mL per included syringe (shake well before using)

CHILDREN'S LIQUID: 160 mg/5 mL per teaspoon (shake well before using)

CHILDREN'S CHEWS OR MELTAWAYS: 80 mg each

JUNIOR-STRENGTH CHEWS OR MELTAWAYS: 160 mg each

ADULT REGULAR-STRENGTH TABLETS: 325 mg each