

The following drugs contain aspirin or other products, which can cause increased bleeding during surgery and should not be taken for 10 days prior to surgery. For relief of minor pain, you MAY take Tylenol (acetaminophen).

Advil	Doan's Regular	Naprelan
Aggrenox	Doan's Extra Strength	Nabumeton
Aleve	Dolobid	Naprosyn
Alka Seltzer	Dristan	Naproxen
Alcohol	Duradyne Tablets	Norgesic Forte
Anacin	Easprin	Omega-3
Anaprox	Ecotrin	Orudis
Ansaid	Eliquis	Oruvail
Arthotec	Empirin	Oxaprozin
Aspirin	Equalgesic Tablets	Pepto Bismol Liquid/Tablets
APC	Etodolac	Percodan
Apixaban	Excedrin	Perantine
BC Tablets or Powders	Feldene	Piroxicam
BC Cold Powder	Fiorinal	Plavix
Brilinta	Fish Oil	Pradaxa
Brufen	Flurbiprofen Sodium	Quagesic
Bufferin	4 Way Cold Tablets	Relafen
Cama Arthritis Pain Reliever	Goody's Headache Powder	Robaxisal
Cataflam	Goody's Headache Tablets	Rufen
Celebrex	Herbal Supplements	Sine Aid
Cephalgesics	Ibuprofen	Soma Compound
Cilostazol	Indomethacine	Sulindac
Clinoril	Indocin	Trandate
Cogesprin	Ketoprofen	Trental
Coricidin	Lodine	Trilisate
Coumadin	Lovenox	Vanquish
Dabigatran	Meclomen	Vitamin E
Darvon	Medipren	Voltaren
Darvon with Aspirin	Meloxicam	Warfarin
Daypro	Midol 200	Wesprin
Diclofenac	Midol PMS Caplets	Xarelto
Diflunisal	Mobic	Zavtrin
Disalcid	Motrin	Zoprin

Before stopping any of these medications, be sure to consult with the physician who ordered them. Some, such as Coumadin (Warfarin) and Plavix, are ordered to prevent or treat serious conditions such as deep venous thrombosis, pulmonary embolisms and other heart problems. This is not an all-inclusive list. If you are unsure if you are taking an aspirin product or an anti-inflammatory, please ask your doctor, nurse or pharmacist.

If you have any questions, please call our office at 630-967-2225.